

Cone-beam Computed Tomography Evaluation of Upper Airway Morphology and its Association with Age, Gender, and Body Mass Index in Healthy Adults



Bayu Trinanda Putra¹ , Inka Saraswati¹ , Menik Priaminiarti¹ , Bramma Kiswanjaya¹ and Hanna H Bachtiar-Iskandar^{1,*}

¹Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Universitas Indonesia, Jakarta, Indonesia

Abstract:

Introduction: This cross-sectional study used Cone-Beam Computed Tomography (CBCT) to evaluate the association between the Upper Airway (UA) volume and demographic risk factors, including age, gender, and Body Mass Index (BMI), in healthy Indonesian adults.

Materials and Methods: A retrospective cross-sectional study was conducted on 85 CBCT datasets (45 males, 40 females; ages 20-40 years). Upper airway segmentation was performed using CS 3D Imaging Software to obtain the measurement parameters, such as total UA volume, minimal cross-sectional area (CSAmin), and lateral and anteroposterior CSAmin dimensions. Inter- and intra-rater reliability were assessed using Intraclass Correlation Coefficients (ICC). Correlation analyses (Spearman's Rho) and group comparisons were used to evaluate associations between airway parameters and age, gender, and Body Mass Index (BMI).

Results: A reliability test for the upper airway measurement parameters was conducted by two examiners over a two-week interval, and the findings indicated that inter-rater reliability was excellent (ICC=0.99), and intra-rater reliability was good (ICC=0.80-0.89). The total UA volume differed significantly by gender (males: $31.7 \pm 7.0 \text{ cm}^3$; females: $27.1 \pm 5.0 \text{ cm}^3$; $p < 0.05$), age group ($p < 0.05$), and BMI category ($p < 0.05$). Weak negative correlations were observed between total UA volume and both gender ($r = -0.327$, $p < 0.05$) and BMI ($r = -0.326$, $p < 0.05$), while age showed no significant correlation. CSAmin and linear dimensions showed no significant correlation with any demographic variable, and all CSAmin values exceeded 110 mm^2 .

Discussion: In this study, upper airway volume was significantly greater in males than in females, consistent with previous findings. Participants with normal BMI exhibited the highest airway volumes, while overweight individuals showed reduced volumes. On the other hand, age did not correlate significantly with upper airway parameters in this 20-40-year-old cohort.

Conclusion: Upper airway volume is significantly associated with gender and BMI, but not with age, in healthy adults aged 20-40 years. Three-dimensional CBCT provides reliable measurements for UA assessment and may support early identification of airway alterations in clinical practice.

Keywords: Cone-beam computed tomography, Upper airway volume, Gender, Body mass index.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Universitas Indonesia, Jakarta, Indonesia; E-mail: hanna.bachtiar@ui.ac.id

Cite as: Putra B, Saraswati I, Priaminiarti M, Kiswanjaya B, Bachtiar-Iskandar H. Cone-beam Computed Tomography Evaluation of Upper Airway Morphology and its Association with Age, Gender, and Body Mass Index in Healthy Adults. Open Dent J, 2026; 20: e187421061521. <http://dx.doi.org/10.2174/01187421061521260710050045>



CrossMark

Received: November 16, 2025

Revised: April 10, 2026

Accepted: May 18, 2026

Published: July 15, 2026



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

The upper airway is a complex anatomical structure

composed of mucosal tissue, musculature, cartilage, and lymphatic elements, and it plays vital roles in respiration and phonation. Alterations in airway morphology may be

caused by infectious or non-infectious pathologies, congenital conditions, or craniofacial anatomical variations [1].

Previous studies have shown that upper airway volume differs across populations and is influenced by several demographic and anatomical factors. For example, studies in India and South Korea reported that males generally have larger upper airway volumes than females [2]. Risk factors associated with upper airway volume include age, gender, and Body Mass Index (BMI) [3-5]. Most earlier investigations relied on two-dimensional cephalometric radiographs, which are limited by magnification errors, distortion, and superimposition of structures, thereby reducing measurement accuracy [6-10].

Cone-Beam Computed Tomography (CBCT) provides three-dimensional visualization of craniofacial structures with high spatial resolution and relatively low radiation dose, making it a valuable tool for airway assessment. CBCT enables accurate segmentation of the nasopharyngeal, oropharyngeal, and hypopharyngeal regions and has been shown to yield highly reproducible airway measurements. Despite these advantages, a study on the efficacy of upper airway measurements using the three-dimensional imaging modality of CBCT on a phantom was conducted by Aboudara *et al.*, who found that the error range between measurements of volume and sagittal area ranged from 0-5% with moderate correlation ($r=0.75$), whereas using the lateral cephalometric radiographic method ranged from 2.5-5.7% [7].

Given the importance of establishing population-specific reference values, this study aimed to evaluate the association between upper airway volume and demographic factors, including age, gender, and BMI, using CBCT images of healthy Indonesian adults aged 20-40 years. Understanding these relationships may

enhance clinical interpretation of airway anatomy and support early identification of individuals at potential risk for airway-related disorders.

2. MATERIALS AND METHODS

This retrospective cross-sectional study was approved by the Ethics Committee of the Faculty of Dentistry, Universitas Indonesia (Number: 68/Ethical Approval/FKGUI/XI/2023). The study was conducted at the Radiology Unit of the dental hospital of Universitas Indonesia over a period of 1 year, from 2023 to 2024. Eighty-five CBCT datasets were collected, and the medical records of patients were reviewed. The following are the inclusion criteria: (1) an adult group with ages ranging from 20-30 years and 31-40 years, and (2) no history of respiratory problems. According to previous studies, the age groups of 20-30 years and 31-40 years were considered stable. Exclusion criteria in this study were data volumes with a Field of View (FOV) of 10 cm x 5 cm or smaller, regardless of coverage of the boundaries of the upper airway. Informed consent was waived by the Ethics Committee of the Faculty of Dentistry, Universitas Indonesia, because this study used retrospectively collected and fully anonymized CBCT datasets, posing minimal risk to participants.

Before determining the boundaries of UA, which are presented in Table 1, six anatomic landmarks that are relevant for upper airway analysis were located, such as (1) sella turcica (S) is the middle point of the sella turcica, (2) Posterior Nasal Spine (PNS) is the most posterior point of the hard palate, (3) anteroposterior C2 is the midinferior point of cervical vertebra C2, (4) superior point of epiglottis is the midsuperior point of epiglottis, (5) base of epiglottis is the inferior point of epiglottis crypt. The boundaries of the upper airway are shown in Fig. (1).

Table 1. The boundaries of the upper airway.

Pharyngeal Borders	Definition
Total Upper Airway	
Anterior boundary	The plane extends through the sella turcica to the PNS, base of the epiglottis, and tip of the epiglottis.
Posterior boundary	The plane extends through the sella turcica to the pharyngeal superior wall parallel to the PNS and the pharyngeal inferior wall parallel to the base of the epiglottis.
Superior boundary	Sella turcica
Inferior boundary	The plane extends through the base of the epiglottis to the inferior posterior wall of the pharynx.
Nasofaring	
Anterior boundary	The plane extends through the sella turcica to the PNS.
Posterior boundary	The plane extends from the sella turcica to the pharyngeal superior wall parallel to the PNS.
Superior boundary	Sella turcica
Inferior boundary	The plane extends through the sella turcica to the pharyngeal superior wall.
Orofaring	
Anterior boundary	The plane extends through the PNS to the tip of the epiglottis.
Posterior boundary	The plane extends through the pharyngeal superior wall to the pharyngeal midposterior wall.
Superior boundary	The plane extends through the PNS to the pharyngeal posterior wall.
Inferior boundary	The plane extends through the tip of the epiglottis to the pharyngeal midposterior wall.
Hipofaring	
Anterior boundary	The plane extends to the base of the epiglottis.
Posterior boundary	The plane extends through the pharyngeal midposterior wall to the pharyngeal inferior wall.
Superior boundary	The plane extends through the tip of the epiglottis to the pharyngeal midposterior wall.
Inferior boundary	The plane extends through the base of the epiglottis to the pharyngeal inferior wall.




Fig. (1). CBCT software (CS 3D Imaging Software 7.0.23 by Carestream Dental, Atlanta) is used to segment the upper airway automatically. **(a)** In the Midsagittal view, the upper airway boundary is marked with a green line as a region of interest, which is formed by the following anatomical landmarks: sella turcica (S), Posterior Nasal Spine (PNS), anteroposterior C2, the midsuperior point of the epiglottis, the base of the epiglottis, and the seed point as a virtual marker region of interest. **(b)** The segmentation results are visible in the Axial view, marked with a blue line. **(c)** The Coronal view is also automatically segmented.

This sample size was calculated using the group mean formula, $N = Z_{\alpha/2}^2 s^2 / d^2$, with an alpha (α) value of 0.05, and a power of 90% was considered. The calculation was based on a previous study conducted by Daya *et.al.* [3]. The determination of six anatomic landmarks and upper airway analysis was performed by two observers, senior residents of oral and maxillofacial radiology who were trained and calibrated using two data sets that were not included in this study.

Eighty-five CBCT data sets were reconstructed, and upper airway segmentation analysis was performed using CBCT software (CS 3D Imaging Software 7.0.23 by Carestream Dental, Atlanta). All CBCT data sets were re-oriented according to the following protocol. In the coronal plane, the most inferior point on the infraorbital margin of both sides lies on the same horizontal plane. In the sagittal plane, the Frankfort plane extends through the ANS (Anterior Nasal Spine) and PNS (Posterior Nasal Spine) points, which are parallel to a horizontal plane. In the axial plane, the plane extending through the ANS to the PNS is perpendicular to the horizontal plane.

The segmentation protocol utilized an airway module where the slice that coincided with the midsagittal plane was selected for the optimal airway view. Anatomic landmark points are connected to form a region of interest in the upper airway, using ruler measurements to delineate the region. After that, 3 “seed points” were added to the center of the region of interest of the nasopharynx, oropharynx, and hypopharynx; these points were used to “expand the airway” (Fig. 1). This method is an automatic segmentation (Fig. 2).

The data collected were sorted and categorized based on the parameters recorded. The upper airway measurement parameters are as follows Total volume upper airway (cm^3), Minimum cross-sectional area (CSAmin) (mm^2), Lateral dimension CSAmin (mm), and Anteroposterior dimension CSAmin (mm).

2.1. Statistical Analysis

The data were analyzed using statistical software. A reliability test was performed by two examiners using the Intraclass Correlation Coefficient (ICC). The null

hypothesis stated that no statistically significant relationship exists between upper airway measurement parameters (total volume upper airway, minimum cross-sectional area (CSAmin), lateral dimension CSAmin, anteroposterior dimension CSAmin) and demographic variables (age, gender, BMI). Data were analyzed using descriptive statistical methods, group comparison tests, and correlation analysis by Spearman's Rho.

3. RESULTS

Inter-rater reliability for upper airway measurement parameters was excellent (ICC = 0.99), while intra-rater reliability ranged from 0.80 to 0.89, indicating good upper airway measurement consistency. A total of 85 CBCT datasets were included in the analysis (45 males, 52.9%; 40 females, 47.1%). The participants were divided into two age groups: 20-30 years ($n = 46$) and 31-40 years ($n = 39$) (Table 2). The BMI categories consisted of underweight (16.5%), normal weight (51.8%), and overweight (31.8%) (Table 3).

The mean total Upper Airway (UA) volume showed significant differences between genders with males presenting higher values than females ($31.7 \pm 7.0 \text{ cm}^3$ vs. $27.1 \pm 5.0 \text{ cm}^3$; $p < 0.05$), likewise, significant differences were also observed across BMI categories ($p < 0.05$), where individuals with normal BMI exhibited the highest UA volumes ($33.2 \pm 5.9 \text{ cm}^3$) compared to underweight ($27.4 \pm 6.4 \text{ cm}^3$) and overweight participants ($24.6 \pm 3.0 \text{ cm}^3$) (Table 4). There are no significant differences in total UA volume between the two age groups ($p > 0.05$). For other airway parameters, including CSAmin, lateral CSAmin, and anteroposterior CSAmin, no statistically significant differences were observed across age group, gender, or BMI.

Spearman's correlation revealed weak but statistically significant negative correlations between total UA volume and both gender ($r = -0.327$, $p < 0.05$) and BMI ($r = -0.326$, $p < 0.05$). Age showed no significant correlation with any airway parameter ($p > 0.05$). All CSAmin values in this study exceeded 110 mm^2 , indicating that participants were generally not at anatomical risk for obstructive sleep apnea based on previously defined CSAmin thresholds (Table 5).

Table 2. Distribution of participants by gender and age group.

Gender	20-30 years, n (%)	31-40 years, n (%)	Total, n (%)
Male	22 (25.9)	23 (28.2)	45 (52.9)
Female	24 (28.2)	16 (18.8)	40 (47.1)
Total	46 (54.1)	39 (45.9)	85 (100.0)

Table 3. Distribution of participants by gender and body mass index (BMI).

Gender	Underweight, n (%)	Normal, n (%)	Overweight, n (%)	Total, n (%)
Male	3 (3.5)	26 (30.6)	16 (18.8)	45 (52.9)
Female	11 (12.9)	18 (21.2)	11 (12.9)	40 (47.1)
Total	14 (16.5)	44 (51.8)	27 (31.8)	85 (100.0)




Fig. (2). 3D segmentation and measuring results (CS 3D Imaging Software 7.0.23 by Carestream Dental, Atlanta). Upper airway segmentation with 3D adjustment (A) and (B), measurement findings for total upper airway volume in cm³, CSAmin in mm², and Lateral and Anteroposterior CSAmin in mm (C).

Table 4. Mean value, standard deviation, and *p*-value for upper airway measurement variables in age, gender, and body mass index groups.

		Total Volume Upper Airway (cm ³)		CSAmin (mm ²)		CSAmin Lateral (mm)		CSAmin Anteroposterior (mm)	
		Mean ± SD	<i>p</i> -value	Mean ± SD Median (Min-Max)	<i>p</i> -value	Mean ± SD Median (Min-Max)	<i>p</i> -value	Mean ± SD	<i>p</i> -value
Age Group	20-30 tahun	30.0 ± 6.2	0.001* ^a	151.3 ± 78.7	0.563 ^a	22.5 ± 7.2	0.960 ^a	9.1 ± 2.7	0.695 ^a
	31-40 tahun	29.0 ± 6.9		141.7 ± 71.8		22.5 ± 6.8		8.9 ± 3.0	
Gender	Male	31.7 ± 7.0	0.001* ^a	138.7 (24.3-324.1)	0.871 ^b	21.8 ± 7.2	0.352 ^a	9.3 ± 3.0	0.357 ^a
	Female	27.1 ± 5.0		130.8 (33.2-365.4)		23.2 ± 6.9		8.7 ± 2.5	
Body Mass Index	Underweight	27.4 ± 6.4	0.000* ^c	157.7 ± 58.2	0.398 ^c	24.4 (10.5-33.5)	0.410 ^d	9.0 ± 2.0	1.000 ^a
	Normal	33.2 ± 5.9		153.4 ± 84.7		22.2 (9.9-45.7)		9.0 ± 3.3	
	overweight	24.6 ± 3.0		130.6 ± 66.0		9.9 (12.3-30.5)		9.0 ± 2.3	

Note: a t-test independent

b Mann-Whitney

c One-way ANOVA

d Kruskal-Wallis

* *p*-value <0.05

Table 5. Spearman-Rho correlation test for upper airway measures (Total upper airway volume, CSAmin, Lateral CSAmin and Anteroposterior CSAmin) with age group, gender and body mass index.

	Total Volume Upper Airway		CSAmin		CSAmin Lateral		CSAmin Anteroposterior	
	Correlation (r)	<i>p</i> -value	Correlation (r)	<i>p</i> -value	Correlation (r)	<i>p</i> -value	Correlation (r)	<i>p</i> -value
Age Group	-0.100	0.364	-0.40	0.713	-0.06	0.960	-0.57	0.606
Gender	-0.327	0.002*	0.18	0.872	0.079	0.471	-0.80	0.468
Body Mass Index	-0.326	0.002*	-0.166	0.128	-0.138	0.207	0.017	0.879

Note: * *p*-value < 0.05.

4. DISCUSSION

The upper airway is a dynamic anatomical structure composed of mucosal, muscular, cartilaginous, and lymphatic components that contribute to respiration and phonation [11, 12]. Alterations in airway morphology are clinically relevant because of reduced airway dimensions [3]. Establishing normative upper airway measurements in healthy populations is critical for identifying deviations that may indicate pathological risk.

The rising prevalence of alteration of the upper airway has been accompanied by advances in diagnostic imaging modalities for evaluating maxillofacial anatomical structures of the upper airway [11]. However, conventional techniques remain limited by factors such as non-linear magnification, image distortion, and superimposition of craniofacial structures, which complicate accurate upper airway assessment [8]. A study by Xing Feng *et al.* comparing total upper airway volume obtained from lateral cephalometry and CBCT in individuals aged < 15 and > 15 years reported a moderate negative correlation ($r = -0.32$). In that study, the Fujioka method, specifically the Adenoidal Nasopharyngeal Ratio

(ANR) derived from lateral cephalometric analysis, was used as a reference for airway volume estimation. Nevertheless, ANR has limited accuracy, as it does not account for physiological variables and is restricted to assessing nasopharyngeal capacity alone [6].

This study found that the volume of the upper airway was significantly greater in males than in females, consistent with previous findings. Elagib *et al.* reported a similar sex-based difference in a South Korean cohort, with males exhibiting larger total airway volumes than females [2]. Di Carlo *et al.* reported that male participants consistently showed larger three-dimensional airway dimensions in orthognathic surgery populations [12]. These differences may be attributed to the larger craniofacial dimensions, greater oropharyngeal soft tissue volume, and differences in fat distribution patterns in males compared to females [8]. This study reinforces these observations by demonstrating a weak but statistically significant correlation between total airway volume and gender.

Body Mass Index (BMI) was also significantly associated with upper airway volume. Participants with

normal BMI exhibited the highest airway volumes, whereas those with overweight showed reduced volumes. This finding aligns with studies indicating that increased adiposity, particularly parapharyngeal fat deposition, may contribute to airway narrowing.

Dastan *et al.* reported comparable results in an Iranian population, showing a significant relationship between airway volume, craniofacial characteristics, and BMI [13]. A weak correlation was found in this study; it remains clinically relevant, as even small volumetric reductions may contribute to airway collapsibility, especially in predisposed individuals.

On the other hand, age did not correlate significantly with upper airway parameters in this 20-40 year old cohort. This is consistent with reports that airway morphology remains relatively stable in early and mid-adulthood and tends to decline only after the fifth decade of life due to progressive soft-tissue laxity and structural remodeling [14]. Since participants in this study were limited to younger adults with presumably stable craniofacial growth, large age-related volumetric differences were not expected.

An important observation from this study is that all CSAm measurements exceeded 110 mm², a threshold previously associated with reduced OSA risk [11]. This supports the classification of the present sample as a healthy, low-risk population. CSAm remains a critical parameter because minimal cross-sectional area rather than total airway volume has been shown to be the most predictive indicator of potential airway obstruction.

The use of CBCT provided high-resolution three-dimensional visualization and reliable segmentation of the upper airway. Both inter-and intra-rater reliability in this study were high, reinforcing the reproducibility of CBCT-based airway assessment [11]. This is consistent with findings by Zimmerman *et al.* and Daya *et al.*, who reported excellent reliability for CBCT-based upper airway measurements when standardized protocols are applied [15]. Despite its advantages, CBCT interpretation must consider patient positioning and breathing phase, as variations in head posture and airway tone can influence volumetric measurements [16].

Overall, this study contributes population-specific normative data for healthy Indonesian adults, a group underrepresented in existing literature. Differences in craniofacial structure across ethnicities make such local data essential for clinical interpretation and accurate risk assessment. Further investigation using larger and more varied samples is required to refine the reference ranges and examine other factors such as craniofacial morphology, soft-tissue characteristics, and functional respiratory assessments.

5. STUDY LIMITATION

This cross-sectional study has several limitations. The retrospective design and relatively small sample size from a single institution may limit the generalizability of the findings. Potential selection bias may also exist because

CBCT scans were initially taken for clinical purposes. Additionally, factors such as patient positioning, breathing phase, and unmeasured anatomical or lifestyle variables could have influenced airway measurements. Future studies with larger and more diverse samples, as well as prospective designs, are needed to validate these results.

CONCLUSION

The findings of this study can be summarized as follows:

- The Upper Airway (UA) volume is significantly associated with gender and BMI, but not with age, in healthy adults aged 20-40 years.
- The factors that influence UA volume include craniofacial dimensions, oropharyngeal soft tissue, and fat distribution.
- Assessment of UA volume should consider head positioning to ensure accurate representation.
- Three-dimensional CBCT provides reliable measurements for UA evaluation.
- CBCT may support early identification of airway alterations in clinical practice.

AUTHORS' CONTRIBUTIONS

The authors confirm contribution to the paper as follows: HB-I, M.P., B.K., B.T.P.: Study Conception and design; B.T.P.: Data Collection; HB-I, M.P., B.K., B.T.P.: Analysis and interpretation of results; HB-I, M.P., B.K., B.T.P.: Draft manuscript. All authors reviewed the results and approved the final version of the manuscript.

LIST OF ABBREVIATIONS

PNS	=	Posterior Nasal Spine
ANS	=	Anterior Nasal Spine
ICC	=	Intraclass Correlation Coefficient
CBCT	=	Cone-Beam Computed Tomography

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study was approved by the Research Ethics Committee of the Faculty of Dentistry, Universitas Indonesia (Number: 68/Ethical Approval/FKGUI/XI/2023).

HUMAN AND ANIMAL RIGHTS

The study was conducted in accordance with the Declaration of Helsinki.

CONSENT FOR PUBLICATION

Informed consent was waived by the Ethics Committee of the Faculty of Dentistry, Universitas Indonesia, because this study used retrospectively collected and fully anonymized CBCT datasets, posing minimal risk to participants.

STANDARDS OF REPORTING

STROBE guidelines were followed.

AVAILABILITY OF DATA AND MATERIALS

The data supporting the findings of the article are available at the Dental Hospital of the Faculty of Dentistry, Universitas Indonesia.

FUNDING

This study was funded by the International Indexed Publication Grants (PUTI) Q3 (Grant NKB.-96/UN2.RST/HKP.05.00/2024) Universitas Indonesia 2024.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Mete A M, Akbudak I H. Functional Anatomy and Physiology of Airway. In: Erbay R H, Ed. Tracheal Intubation. London: IntechOpen 2018.
<http://dx.doi.org/10.5772/intechopen.77037>
- [2] Elagib T, Kyung H, Hung BQ, Hong M. Assessment of pharyngeal airway in Korean adolescents according to skeletal pattern, sex, and cervical vertebral maturation: A cross-sectional CBCT study. Korean J Orthod 2022; 52(5): 345-53.
<http://dx.doi.org/10.4041/kjod21.267>
- [3] Daya K J, Kajol M T, Vikram G, Abhijeet V P, Sunaina M S, Pranali B B. Assessment of interobserver reliability in localization of hard and soft tissue landmarks and three-dimensional measurements of the upper airway based on these landmarks using CBCT images. IP Int J Maxillofac Imaging 2021; 7(2): 54-60.
<http://dx.doi.org/10.18231/ijmi.2021.011>
- [4] Zinsly SDR, Moraes LCD, Moura PD, Ursi W. Open-access Evaluation of the pharyngeal airway space using cone-beam computed tomography. Dental Press J Orthod 2010; 15(5): 150-8.
<http://dx.doi.org/10.1590/S2176-94512010000500018>
- [5] Emsaeili F, Sadrhaghghi A, Sadeghi-Shabestari M, Nastarin P, Niknafs A. Comparison of superior airway dimensions and cephalometric anatomic landmarks between 8-12-year-old children with obstructive sleep apnea and healthy children using CBCT images. J dent res dent clin dent prospects 2022; 16(1): 18-23.
<http://dx.doi.org/10.34172/joddd.2022.003> PMID: 35936930
- [6] Feng X, Li G, Qu Z, Liu L, Näsström K, Shi X. Comparative analysis of upper airway volume with lateral cephalograms and cone-beam computed tomography. Am j orthod dentofac orthop : off publ Am Assoc Orthod const soc Am Board Orthod 2015; 147(2): 197-204.
<http://dx.doi.org/10.1016/j.ajodo.2014.10.025> PMID: 25636553
- [7] Aboudara C, Nielsen I, Huang JC, Maki K, Miller AJ, Hatcher D. Comparison of airway space with conventional lateral headfilms and 3-dimensional reconstruction from cone-beam computed tomography. Am j orthod dentofac orthop : off publ Am Assoc Orthod const soc Am Board Orthod 2009; 135(4): 468-79.
<http://dx.doi.org/10.1016/j.ajodo.2007.04.043> PMID: 19361733
- [8] Buchanan A, Cohen R, Looney S, Kalathingall S, De Rossi S. Cone-beam CT analysis of patients with obstructive sleep apnea compared to normal controls. Imaging sci dent 2016; 46(1): 9-16.
<http://dx.doi.org/10.5624/isd.2016.46.1.9> PMID: 27051634
- [9] Bajrovic N, Nakas E, Dzemic V, Tiro A. The Link Between Obstructive Sleep Apnea and Orthodontic Anomalies in Obese Adult Population. Mater socio-med 2021; 33(1): 65-9.
<http://dx.doi.org/10.5455/msm.2021.33.65-69> PMID: 34012354
- [10] Benjafield AV, Ayas NT, Eastwood PR, et al. Estimation of the global prevalence and burden of obstructive sleep apnoea: a literature-based analysis. Lancet Respir Med 2019; 7(8): 687-98.
[http://dx.doi.org/10.1016/s2213-2600\(19\)30198-5](http://dx.doi.org/10.1016/s2213-2600(19)30198-5)
- [11] Chaudhry U, Cohen JR, Al-Samawi Y. Use of cone beam computed tomography imaging for airway measurement to predict obstructive sleep apnea. CRANIO® 2020; 40(5): 418-24.
<http://dx.doi.org/10.1080/08869634.2020.1765602>
- [12] Di Carlo G, Gurani SF, Pinholt EM, Cattaneo PM. A new simple three-dimensional method to characterize upper airway in orthognathic surgery patient. Dento maxillo facial radiol 2017; 46(8): 20170042.
<http://dx.doi.org/10.1259/dmfr.20170042> PMID: 28749703
- [13] Dastan F, Ghaffari H, Shishvan HH, Zareiyani M, Akhlaghian M, Shahab S. Correlation between the upper airway volume and the hyoid bone position, palatal depth, nasal septum deviation, and concha bullosa in different types of malocclusion: A retrospective cone-beam computed tomography study. Dent med probl 2021; 58(4): 509-14.
<http://dx.doi.org/10.17219/dmp/130099> PMID: 34850611
- [14] Eow PY, Lin KY, Kohli S, Math SY. Cone-beam computed tomography assessment of upper airway dimensions in patients at risk of obstructive sleep apnea identified using STOP-Bang scores. Imaging Sci Dent 2021; 51(4): 439.
<http://dx.doi.org/10.5624/isd.20210193>
- [15] Zimmerman JN, Vora SR, Pliska BT. Reliability of upper airway assessment using CBCT. Eur J Orthod 2018; 41(1): 101-8.
<http://dx.doi.org/10.1093/ejo/cjy058>
- [16] Anandarajah S, Abdalla Y, Dudhia R, Sonnesen L. Proposal of new upper airway margins in children assessed by CBCT. Dentomaxillofac Radiol 2015; 44(7): 20140438.
<http://dx.doi.org/10.1259/dmfr.20140438>

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.